

# Initial Estimate of the $^{237}\text{U}(\text{n},\text{f})$ Cross Section for $0.1 < E_{\text{n}}(\text{MeV}) < 20$

*W. Younes, H. C. Britt, J. A. Becker, J. B. Wilhelmy*

**June 27, 2003**

**U.S. Department of Energy**

Lawrence  
Livermore  
National  
Laboratory

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

# Initial estimate of the $^{237}\text{U}(n, f)$ cross section for $0.1 < E_n \text{ (MeV)} \leq 20$

W. Younes,<sup>†</sup> H. C. Britt, and J. A. Becker

*Lawrence Livermore National Laboratory, Livermore, CA 94551*

J. B. Wilhelmy

*Los Alamos National Laboratory, Los Alamos, NM 87545*

(Dated: June 27, 2003)

In response to a request for a  $^{237}\text{U}(n, f)$  cross section evaluation up to  $E_n = 20$  MeV, we have married a data set from our previous reliable estimate [1–3] of the cross section up to  $E_n = 2.5$  MeV, to an estimate of the remaining cross section up to  $E_n = 20$  MeV, deduced from simple physics arguments. This straw-man, work-in-progress estimate of the  $^{237}\text{U}(n, f)$  cross section is intended to be used in sensitivity-test comparisons to other evaluations of the cross section (e.g., ENDF/B-VI [4] and ENDL [5]). The simple approach used in this work to generate a consistent cross section up to  $E_n = 20$  MeV is validated using the well-known  $^{235}\text{U}(n, f)$  cross section as a test case (see Fig. 1). The corresponding estimate of the  $^{237}\text{U}(n, f)$  cross section is plotted in Fig. 2 and listed in Table I.

In a separate report [6], our  $^{237}\text{U}(n, f)$  cross section estimate in the  $E_n = 0.1 - 2.5$ -MeV range, which is deduced from surrogate-reaction data, has been folded with a “tamped flattop” neutron flux, and normalized to the  $^{235}\text{U}(n, f)$  integral cross section. The value of this ratio agrees within experimental uncertainties with the measured value [7]. The same integral cross-section ratio extracted using the ENDL evaluations for the  $^{237}\text{U}(n, f)$  and  $^{235}\text{U}(n, f)$  cross sections exceeds our result and the measured value by 75%.

A more sophisticated approach, taking proper account of level densities and pre-equilibrium processes, is planned to produce a reliable and robust  $^{237}\text{U}(n, f)$  cross section estimate up to  $E_n = 20$  MeV.

## The $^{235}\text{U}(n, f)$ cross section

The  $^{235}\text{U}(n, f)$  cross section plotted in Fig. 1 was estimated using the formula

$$\begin{aligned} \sigma_{(n,f)}(E_n) = & \sigma_{(n,f)}^{(1)}(E_n) \\ & + \sigma_{(n,f)}^{(2)}(E_n) \times \left[ 1 - \frac{\sigma_{(n,f)}^{(1)}(E_n)}{\sigma_{CN}(E_n)} \right] \\ & \times \{1 - P_{pe}(E_n) \times [1 - P_{pe,f}(E_n)]\}, \quad (1) \end{aligned}$$

where  $\sigma_{(n,f)}^{(1)}(E_n)$  is the first-chance fission cross section, consisting of our previous data continued by a linear extrapolation up to  $E_n = 20$  MeV, and  $\sigma_{(n,f)}^{(2)}(E_n)$  is the second-chance fission cross section. The second-chance fission cross section is reduced, 1) by a factor that removes that fraction of the compound-nucleus cross section lost to first-chance fission, and 2) by a second factor to remove those pre-equilibrium events (characterized by the probability  $P_{pe}$ ) that leave the residual nucleus with an excitation energy below the fission barrier (characterized by  $1 - P_{pe,f}$ ).

In the range  $0.1 < E_n \text{ (MeV)} \leq 2.2$ , we have used our previous results [1–3] for the first-chance fission. We have extended these results to  $E_n = 20$  MeV by assuming a linear dependence of the first-chance fission cross section on the incident neutron energy. The slope of the line is taken to be the same as in the ENDF/B-VI evaluation [4] over the  $E_n = 2 - 5.5$ -MeV range. The intercept is fixed by matching the line at  $E_n = 2.2$  MeV to the average  $^{235}\text{U}(n, f)$  cross section for  $E_n = 1 - 2.5$  MeV (1.23 barns) deduced in our previous work [1].

The second chance fission, which corresponds to fission of the  $^{235}\text{U}$  compound nucleus, is simulated by using the well-known  $^{234}\text{U}(n, f)$  cross section [4], shifted in neutron energy to match the excitation energy that would be populated in the  $^{235}\text{U}(n, n')$  reaction:

$$\begin{aligned} \sigma_{(n,f)}^{(2)}(E_n) \equiv & \sigma_{(n,f)}^{(2)}(A, E_n) \\ = & \sigma_{(n,f)}(A - 1, E_n - B_n(A) - \Delta E), \quad (2) \end{aligned}$$

where we’ve explicitly written the dependence on the target nucleus, represented by its atomic mass  $A$ . The quan-




FIG. 1: Estimated  $^{235}\text{U}(n, f)$  cross section, compared to the ENDF/B-VI evaluation.

tity  $\Delta E$  represents the kinetic energy lost by the neutron emitted in the  $^{235}\text{U}(n, n')$  reaction, and the binding energy is  $B_n(235) = 5.298$  MeV. In practice, the neutron energy shift  $-B_n(A) - \Delta E$  was adjusted to produce the best fit to the ENDF/B-VI  $^{235}\text{U}(n, f)$  cross section, from which a value  $\Delta E \approx 0.70$  MeV was deduced. The experimental  $^{234}\text{U}(n, f)$  cross section used in Eq. (2) includes an implicit contribution from third-chance fission, but no correction has been made for pre-equilibrium effects in the  $^{235}\text{U}(n, 2nf)$  channel.

The probability of a pre-equilibrium neutron emission  $P_{pe}(E_n)$  was calculated using the code DDHMS [8]. The fraction of pre-equilibrium leaving the residual nucleus  $^{235}\text{U}$  with an excitation energy above the inner fission barrier height  $E_A = 6.00$  MeV [9] was deduced from the centroid  $\overline{E}_x^{(pe)}$  and standard deviation  $\sigma_{pe}$  of the population after pre-equilibrium emission predicted by DDHMS, by assuming a Gaussian population distribution,

$$P_{pe,f}(E_n) = \frac{1}{\sqrt{2\pi}\sigma_{pe}} \int_{E_A}^{\infty} dE_x e^{-\frac{(\overline{E}_x^{(pe)} - E_x)^2}{2\sigma_{pe}^2}}. \quad (3)$$

The estimated  $^{235}\text{U}(n, f)$  cross section in Fig. 1 is in good agreement with the ENDF/B-VI evaluation, with localized deviations of 20% or less.

### The $^{237}\text{U}(n, f)$ cross section

The procedure tested with the  $^{235}\text{U}(n, f)$  cross section was applied to the  $^{237}\text{U}(n, f)$  case. Our previous  $(n, f)$  results [1–3] were used for  $0.1 < E_n$  (MeV)  $\leq 2.5$ . These first-chance fission results were extrapolated with a linear function of  $E_n$  whose slope was taken to be the same as in the ENDF/B-VI evaluation for the  $^{237}\text{U}(n, f)$  cross section in the  $E_n = 2 - 5.5$ -MeV range. The linear extrapolation is flatter than in the  $^{235}\text{U}(n, f)$  case. To lowest order, the slope of the first-chance fission cross section is a function of the difference between the barrier height and the neutron binding energy [10]. That energy difference is larger in the case of  $^{237}\text{U}(n, f)$  compared to  $^{235}\text{U}(n, f)$ , and simple statistical arguments confirm that the first-chance fission cross section should be flatter for the  $^{237}\text{U}(n, f)$  reaction [10].

The second-chance fission was simulated using the well-known  $^{236}\text{U}(n, f)$  cross section [4] with a neutron energy shift,  $-B_n(237) - \Delta E$ , and using the value  $\Delta E = 0.70$  MeV determined in the  $^{235}\text{U}(n, f)$  case. Corrections for pre-equilibrium neutron emission were applied as in Eq. (1), using the code DDHMS. The inner fission barrier height  $E_A = 6.30$  MeV [9] was used in Eq. (3).

The estimated  $^{237}\text{U}(n, f)$  cross section is plotted in Fig. 2, and given explicitly in Table I. The ENDF/B-VI and ENDL [5] evaluations are in good agreement with each other above  $E_n \approx 5$  MeV, but differ significantly at lower neutron energies. The cross section estimated in this work does not agree well with either evaluation, especially for  $E_n \leq 5$  MeV.

TABLE I: Estimated  $^{237}\text{U}(n, f)$  cross section.

| $E_n$ (MeV) | $\sigma_{(n,f)}(E_n)$ (b) |
|-------------|---------------------------|
| 0.15        | 0.62                      |
| 0.20        | 0.58                      |
| 0.25        | 0.57                      |
| 0.30        | 0.55                      |
| 0.35        | 0.56                      |
| 0.40        | 0.54                      |
| 0.45        | 0.53                      |
| 0.50        | 0.54                      |
| 0.55        | 0.55                      |
| 0.60        | 0.53                      |
| 0.65        | 0.54                      |
| 0.70        | 0.55                      |
| 0.75        | 0.55                      |
| 0.80        | 0.57                      |
| 0.85        | 0.51                      |
| 0.90        | 0.51                      |
| 0.95        | 0.49                      |
| 1.00        | 0.47                      |
| 1.05        | 0.48                      |
| 1.10        | 0.45                      |
| 1.15        | 0.48                      |
| 1.20        | 0.48                      |




FIG. 2: Estimated  $^{237}\text{U}(n, f)$  cross section, compared to the ENDF/B-VI and ENDL evaluations.

TABLE I: (Continued).

| $E_n$ (MeV) | $\sigma_{(n,f)}(E_n)$ (b) |
|-------------|---------------------------|
| 1.25        | 0.48                      |
| 1.30        | 0.47                      |
| 1.35        | 0.48                      |
| 1.40        | 0.46                      |
| 1.45        | 0.43                      |
| 1.50        | 0.43                      |
| 1.55        | 0.48                      |
| 1.60        | 0.45                      |
| 1.70        | 0.46                      |
| 1.75        | 0.46                      |
| 1.80        | 0.49                      |
| 1.85        | 0.50                      |
| 1.90        | 0.52                      |
| 1.95        | 0.52                      |
| 2.00        | 0.54                      |
| 2.04        | 0.50                      |
| 2.09        | 0.52                      |
| 2.14        | 0.52                      |
| 2.19        | 0.52                      |
| 2.24        | 0.53                      |
| 2.29        | 0.51                      |
| 2.33        | 0.52                      |
| 2.38        | 0.51                      |
| 2.50        | 0.49                      |

TABLE I: (Continued).

| $E_n$ (MeV) | $\sigma_{(n,f)}(E_n)$ (b) |
|-------------|---------------------------|
| 3.00        | 0.49                      |
| 3.50        | 0.48                      |
| 4.00        | 0.48                      |
| 4.50        | 0.48                      |
| 5.00        | 0.47                      |
| 5.50        | 0.47                      |
| 6.00        | 0.47                      |
| 6.50        | 0.50                      |
| 7.00        | 0.92                      |
| 7.50        | 1.00                      |
| 8.00        | 1.10                      |
| 8.50        | 1.11                      |
| 9.00        | 1.11                      |
| 9.50        | 1.11                      |
| 10.00       | 1.10                      |
| 10.50       | 1.09                      |
| 11.00       | 1.09                      |
| 11.50       | 1.13                      |
| 12.00       | 1.27                      |
| 12.50       | 1.43                      |
| 13.00       | 1.57                      |
| 13.50       | 1.62                      |
| 14.00       | 1.64                      |
| 14.50       | 1.69                      |
| 15.00       | 1.70                      |
| 15.50       | 1.67                      |
| 16.00       | 1.63                      |
| 16.50       | 1.61                      |
| 17.00       | 1.62                      |
| 17.50       | 1.64                      |
| 18.00       | 1.65                      |
| 18.50       | 1.66                      |
| 19.00       | 1.69                      |
| 19.50       | 1.73                      |
| 20.00       | 1.78                      |

## REFERENCES

- <sup>†</sup> Electronic address: [younes@llnl.gov](mailto:younes@llnl.gov)
- [1] W. Younes and H. C. Britt (2003), submitted to Phys. Rev. C.
  - [2] W. Younes, H. C. Britt, and J. B. Wilhelmy, *The  $^{237}\text{U}(n, f)$  cross section*, Tech. Rep. UCRL-ID-152621, LLNL (2003).
  - [3] W. Younes and H. C. Britt, *Tables of neutron-induced fission cross sections for various Pu, U, and Th isotopes, deduced from measured fission probabilities*, Tech. Rep. UCRL-ID-152906, LLNL (2003).
  - [4] Cross Section Evaluation Working Group, *ENDF/B-VI Summary Documentation*, Report **BNL-NCS-17541 (ENDF-201)** (1991), edited by P. F. Rose, National Nuclear Data Center, Brookhaven Laboratory, Upton, NY, USA.
  - [5] *Evaluated neutron data library*, [www-ndg.llnl.gov](http://www-ndg.llnl.gov).
  - [6] W. Younes and H. C. Britt, *Neutron fission of  $^{235,237,239}\text{U}$  and  $^{241,243}\text{Pu}$ : cross sections, integral cross sections, and cross sections on excited states*, Tech. Rep., LLNL (2003), manuscript in preparation.
  - [7] D. W. Barr, private communication.
  - [8] M. B. Chadwick, private communication.
  - [9] H. C. Britt, in *Proceedings of the Fourth International Symposium on the Physics and Chemistry of Fission*, Julich, West Germany (International Atomic Energy Agency, Vienna, 1979), vol. 1, p. 3.
  - [10] P. M. Endt and P. B. Smith, eds., *Nuclear Reactions* (North-Holland Publishing Company, Amsterdam, 1962), vol. 2, chap. 2.